Author index

Afify, E.A.

Turnover of μ-opioid receptors in neuroblastoma cells (106) 83 Akil, H., see Isgor, C. (106) 30 Appleby, V.J., see Lee, C.-J. (106) 151 Aschner, M., see Shanker, G. (106) 1

Banno, Y., see Nakagawa, M. (106) 12
Bogoch, I.I., see Dyck, R.H. (106) 22
Bond, B.C., Virley, D.J., Cairns, N.J., Hunter, A.J., Moore, G.B.T., Moss, S.J., Mudge, A.W., Walsh, F.S., Jazin, E. and Preece, P.
The quantification of gene expression in an animal model of brain.

The quantification of gene expression in an animal model of brain ischaemia using TaqMan™ real-time RT-PCR (106) 101

Cairns, N.J., see Bond, B.C. (106) 101 Chan, W.-I., see Lee, C.-J. (106) 151 Charles, K.J., see Robbins, M.J. (106) 136 Cheng, Y.-C., see Lee, C.-J. (106) 151 Cheung, M., see Lee, C.-J. (106) 151

Dib-Hajj, S., see Renganathan, M. (106) 70 Dyck, R.H., Bogoch, I.I., Marks, A., Melvin, N.R. and Teskey, G.C. Enhanced epileptogenesis in \$100B knockout mice (106) 22

Harrison, D.C., see Robbins, M.J. (106) 136 Hibberd, C., see Yau, J.L.W. (106) 117 Huang, G.C., see Isgor, C. (106) 30 Hunter, A.J., see Bond, B.C. (106) 101

Isgor, C., Huang, G.C., Akil, H. and Watson, S.J.

Correlation of estrogen β-receptor messenger RNA with endogenous levels of plasma estradiol and progesterone in the female rat hypothalamus, the bed nucleus of stria terminalis and the medial amygdala (106) 30

Ishikawa, Y., Katoh, H., Nakamura, K., Mori, K. and Negishi, M. Developmental changes in expression of small GTPase RhoG mRNA in the rat brain (106) 145

Iwama, T., see Nakagawa, M. (106) 12

Jazin, E., see Bond, B.C. (106) 101

Kaku, Y., see Nakagawa, M. (106) 12

Katoh, H., see Ishikawa, Y. (106) 145
Kerokoski, P., Suuronen, T., Salminen, A.,
Soininen, H. and Pirttilä, T.
Influence of phosphorylation of p35,
an activator of cyclin-dependent
kinase 5 (cdk5), on the proteolysis of
p35 (106) 50

Kubo, T., Nishimura, S. and Oda, T. Amyloid β-peptide alters the distribution of early endosomes and inhibits phosphorylation of Akt in the presence of 3-(4,5-dimethylthiazol-2yl)-2,5-diphenyltetrazolium bromide (MTT) (106) 94

Lee, C.-J., Chan, W.-I., Cheung, M., Cheng, Y.-C., Appleby, V.J., Orme, A.T. and Scotting, P.J. CIC, a member of a novel subfamily of the HMG-box superfamily, is transiently expressed in developing granule neurons (106) 151

Marks, A., see Dyck, R.H. (106) 22 Melvin, N.R., see Dyck, R.H. (106) 22 Moore, G.B.T., see Bond, B.C. (106) 101 Mori, K., see Ishikawa, Y. (106) 145 Moss, S.J., see Bond, B.C. (106) 101 Mudge, A.W., see Bond, B.C. (106) 101 Mutkus, L.A., see Shanker, G. (106) 1

Nagane, Y., see Utsugisawa, K. (106) 88
Nakagawa, M., Nakashima, S., Banno, Y.,
Yamada, J., Sawada, M., Yoshimura, S.i., Kaku, Y., Iwama, T., Shinoda, J. and Sakai, N.
Increased expression and secretion of r-Gsp protein, rat counterpart of complement C1s precursor, during cyclic AMP-induced differentiation in rat C6 glioma cells (106) 12

Nakamura, K., see Ishikawa, Y. (106) 145 Nakashima, S., see Nakagawa, M. (106) 12 Negishi, M., see Ishikawa, Y. (106) 145 Nishimura, S., see Kubo, T. (106) 94 Noble, J., see Yau, J.L.W. (106) 117

Obara, D., see Utsugisawa, K. (106) 88 Oda, T., see Kubo, T. (106) 94 Orme, A.T., see Lee, C.-J. (106) 151

Pangalos, M.N., see Robbins, M.J. (106) 136 Pirttilä, T., see Kerokoski, P. (106) 50 Preece, P., see Bond, B.C. (106) 101

Quirin-Stricker, C., see Robert, I. (106) 124

Renganathan, M., Dib-Hajj, S. and Waxman, S.G. Na,1.5 underlies the 'third TTX-R sodium current' in rat small DRG

neurons (106) 70

Robbins, M.J., Charles, K.J., Harrison, D.C. and Pangalos, M.N.

Localisation of the GPRC5B receptor in the rat brain and spinal cord (106)

Robert, I., Sutter, A. and Quirin-Stricker, C. Synergistic activation of the human choline acetyltransferase gene by c-Myb and C/EBPB (106) 124

Sakai, N., see Nakagawa, M. (106) 12 Salminen, A., see Kerokoski, P. (106) 50 Sawada, M., see Nakagawa, M. (106) 12 Scotting, P.J., see Lee, C.-J. (106) 151 Seckl, J.R., see Yau, J.L.W. (106) 117 Shanker, G., Mutkus, L.A., Walker, S.J. and Aschner, M.

Methylmercury enhances arachidonic acid release and cytosolic phospholipase A₂ expression in primary cultures of neonatal astrocytes (106) 1

Shinoda, J., see Nakagawa, M. (106) 12 Sogayar, M.C., see Vedoy, C.G. (106) 57 Soininen, H., see Kerokoski, P. (106) 50 Sutter, A., see Robert, I. (106) 124 Suuronen, T., see Kerokoski, P. (106) 50

Teskey, G.C., see Dyck, R.H. (106) 22 Tohgi, H., see Utsugisawa, K. (106) 88

Utsugisawa, K., Nagane, Y., Obara, D. and Tohgi, H. Over-expression of α7 nicotinic acetylcholine receptor induces sustained ERK phosphorylation and N-cadherin expression in PC12 cells (106) 88

Vedoy, C.G. and Sogayar, M.C.
Isolation and characterization of genes
associated with the anti-tumor activity
of glucocorticoids (106) 57
Virley, D.J., see Bond, B.C. (106) 101

Walker, S.J., see Shanker, G. (106) 1 Walsh, F.S., see Bond, B.C. (106) 101 Waltereit, R. and Weller, M.

The role of caspases 9 and 9-short (9S) in death ligand- and druginduced apoptosis in human astrocytoma cells (106) 42 Watson, S.J., see Isgor, C. (106) 30 Waxman, S.G., see Renganathan, M. (106) 70 Weller, M., see Waltereit, R. (106) 42

Yamada, J., see Nakagawa, M. (106) 12 Yau, J.L.W., Hibberd, C., Noble, J. and Seckl, J.R. The effect of chronic fluoxetine treatment on brain corticosteroid receptor mRNA expression and spatial memory in young and aged rats (106) 117

Yoshimura, S.-i., see Nakagawa, M. (106) 12

